



haustorium

Parasitic
Plants
Newsletter

Number 16
March 1986

Official Organ of the
International Parasitic
Seed Plant Research Group

HELP! HAUSTORIUM IS A PARASITE WITHOUT A HOST!

This could be the last issue of our newsletter as, once again, we are without a sponsor. This issue is being printed and mailed through the generosity of the INTSORMIL office but this is the last issue they can support. Can any one help?

STRIGA SPECIES IN ETHIOPIA

On a recent visit to Ethiopia it was confirmed that *Striga hermonthica* is continuing to spread and is now found up to an elevation of 2300M in some areas. Occurrences on some new farms at low altitude are believed to have arisen from the abundant natural infestation in the native savannah, apparently parasitic mainly of species on *Setaria*.

The most striking species was *S. latericea* which is known from native vegetation in a number of districts in Ethiopia and other parts of East Africa but occurs as a persistent localized problem on the Meta Hare Sugar Corporation farm in the Awash valley. It is as tall as *S. hermonthica* but has broader leaves and a dense covering of fine hairs and spikes of brick-red flowers up to 2 cm long.

Close examination of this population showed that it is perennial with a system of rhizomes several mm thick from

which adventitious buds produce aerial shoots. The aerial parts also have a perennial habit with new shoots arising from the lower nodes after much of the stem has matured and died. The parasite is apparently slow to establish and is not normally noticed in the first year after planting sugar cane, but is seen as spreading patches in ratoons. These patches grow up to several meters across and persist for many years and even re-appear in the same place after the ratoon is destroyed and the crop replanted. New infestations are not often noted and it appears that it spreads mainly by vegetative reproduction. Very little seed was being set due to heavy infestation by a pollen eating larva.

C Parker

WHAT EVER HAPPENED TO THE INDEX OF PARASITIC SEED PLANT WORKERS?

We still plan to produce such an index but production has been delayed due to a change in the way HAUSTORIUM is prepared and, at present, a lack of a sponsor! It is still not too late to send your forms to L J Musselman.

A TUBEROUS HAUSTORIUM OF THONNINGIA SANGUINEA (BALANOPHORACEAE) GROWING ON HEVEA BRASILIENSIS.

In 1985 a tuber 8 cm in diameter was sent to Kew from Cameroun where it was collected by Mr P G S Hall of the Natural Resources

Department, Commonwealth Development Corporation. It was said that *Thonningia sanguinea* was conspicuous as red rosettes on the ground in a rubber plantation. As far as we know such tubers have never been reported from this species and examination of herbarium material at Kew provided nothing like it. Although there was no reason to doubt its identity as photographs of the flowers were provided, no inflorescence was attached so confirmation was sought by anatomical study. The woody root to which the tuber was attached proved to be rubber (*Hevea brasiliensis*) while the tuber consists of parenchymatous ground tissue with islands of vascular tissue pursuing an irregular course and some sclereids. This is anatomically similar to the only reference slide at Kew of another member of the same family, *Langsdorffia papuana* from New Guinea, which is good evidence that the tuber consists of *Thonningia* rather than *Hevea* tissue.

It would be interesting to know whether such a tuber is frequent and whether or not it occurs on host plants other than rubber. According to the Anatomy of the Dicots, tuberous rhizomes in the Balanophoraceae range in size from a small nut to a human head. *Striga gesnerioides* also forms a tuber-like structure of some size but only when the host is an arborescent species of *Euphorbia*, which like *Hevea*, is a latex producing member of the Euphorbiaceae. Is

there an analagous function in these two parasites from totally unrelated families each producing tuberous haustoria? (See figure on page 4).

F N Hepper and P Gasson, Royal Botanic Gardens, Kew.

FOURTH SYMPOSIUM ON PARASITIC WEEDS, SUMMER 1987.

Plans are proceeding for our next IPSPRG meeting which will be held in Germany during the summer of 1987 at the Phillips University in Marburg.

The actual date of the meeting has not yet been decided but will be either before or after the Botanical Congress to be held in Berlin 24 July to 1 August 1987.

MEDICINAL USES OF A MEMBER OF THE BALANOPHORACEAE IN SOMALIA

During a recent collecting trip in the Middle Juba Region of Somalia, we encountered a preparation in the local markets sold as a cure for diarrhea and menstrual disorders called in Somali, *dinsi*. Because of its resemblance to tartous (a member of the Hydnoraceae used in other parts of Africa as a medicine for the same ailments), we attempted to locate the source of the *dinsi*. After some consultation with local people we found that what was being sold was the dried and broken pieces of a member of the *Balanophoraceae*. The plant has not yet been identified but it does not resemble plants of the genus *Balanophora* and could be a species of *Chlamydomyrtum* or a related genus. Further work is under way to determine the plant and other uses as well as some information

on the chemical makeup of the medicine.

Cistanche is also known as *dinsi* in the same area and we were led by a nomad to a stand of *Cistanche* and told it was *dinsi*. However, the material being sold in the market definitely was not *Cistanche*.

Aweys Yusef and L J Musselman

EFFECT OF FERTILIZER ON STRIGA COUNT IN WHEAT

An experiment on the long range effect of continuous cropping and manuring on Jowar wheat rotation is in progress at the Agricultural Research Station of the University of Agricultural Sciences at Sirguppa in the Tunga Bhadra Project area. The soil is a vertisol and the experiment has been in progress since 1977. The treatments consist of all combinations of three levels of N (40, 80, and 120 kg N/ha), three levels of P₂O₅ (0, 40, and 80 kg/ha) and two levels of K₂O (0 and 40 kg/ha). The experiment is laid out in a 3² x 2 partially unfounded design with four replications.

The crop of Hy. jowar-CSH-5 was sown on 8-7-1985 with a spacing of 40 cm between rows and 10 cm between plants within a row. Counts of *Striga asiatica* were recorded treatment-wise at 70 days after sowing. The data on weed counts and visual observation indicates that the weed population is low in the plots receiving higher doses of N. The effect of P₂O₅ levels and K₂O level did not show any influence on the *Striga* count. The data indicates that the intensity of *Striga* is greater in N poor soils.

M M Hosmani, V Jagannath, K M S Sharma, University of

Agricultural Sciences, Shimoga, India.

LITERATURE

The underground sorcery of witchweed. Discovery. December 1985. (This is a popular account of a very significant and as yet unpublished discovery of the "chemical radar" of *Striga asiatica*. J L Riopel and Lynn have found that the parasite sends a message to the host which in turn tells the parasite to produce a haustorium).

Yatskievych, G, Zavada, I 1984. Pollen morphology of Lennoaceae. Pollen et Spores 26: 19-30. (Pollen structure supports the concept that the North American family of holoparasites consists of only *fastidiosa* species.)

Iranshar, M. 1983. New record of cuscuta (Cuscutaceae) from Iran. Iranian Journal of Botany 2(1): 9-12.

(Not surprisingly, *C. campestris* is now known from Iran. Also noted for the first time is *C. lehmanniana*. There are figures of the species and a map of their distribution.)

International Institute of Agriculture. 1985. IITA Research Highlights for 1984. (Two articles on *Striga*, one describing a survey of *Striga* species in maize in Nigeria and suggesting that *S. aspera* is much more important than previously thought and also that *S. forbesii* and a yellow flowered form of *S. asiatica* are of more local importance. Another species, perhaps *S. passargei*, has also been noted on maize. A second article describes the discovery

of resistance to S. gesnerioides in the cowpea variety %vita-2 and the transfer of the resistance into varieties with insect resistance.).

Vasudeva Rao, M J . 1485. Techniques for screening sorghums for resistance to Striga. Information Bulletin 20, ICRISAT. (An extremely well-illustrated and clear guide to a range of techniques for the study of Striga from laboratory to field which will be useful not only to the researcher on crop resistance but to those working on other approaches also. This forms a chapter in the forthcoming volume Striga Biology and control to be published by CRC Press in 1986.)

Kuijt, J, Bray, D, Olson, R. 1985. Anatomy and ultrastructure of the endophytic system of Pilostyles thurberi (Rafflesiaceae). Canadian Journal of Botany 63: 1231-1240. (Three cell types are described from the cortical strands one of which is considered to be a sieve element although vestigial.)

Nassib, A M , Hussein, A H A, El Rayes, F M. 1985. Effect of variety, chemical control, sowing date and tillage on Orobanch spp infestation and faba bean yield. Fabis Newsletter 10:11-15. (A useful summary of a wide ranging series of studies on O. crenata in Egypt.)

Yatskievych, G. 1985. Notes on the biology of the Lennoacme. Cactus and Succulent Journal (U.S.) 57: 73-79. (A well illustrated, in color, and interesting account of this fascinating family.)

Scrophulariaceae Research Newsletter 1(2). (This may be of interest to HAUSTORIUM readers who are work with parasitic Scrophulariaceae. Most of the newsletter is concerned with non-parasitic species but there is also a helpful review of literature which covers the entire family. You can obtain the newsletter by writing: K Barringer, Field Museum of Natural History, Chicago, Illinois 60605.)

Olson, A R, Kuijt, J. 1985. Sieve elements in the morphologically reduced mistletoe Viscum minimum Harvey (Viscaceae). American Journal of Botany 72: 1220-1224. (This miniature mistletoe with shoots only 2-3 mm long on Euphorbia horinda is shown to have functional phloem elements, unlike some other reduced mistletoe species.)

Clay, K, Dement, D, Rejmanek, M 1985. Experimental evidence for host races in mistletoe (Phoradendron tomentosum). American Journal of Botany 72: 1225-1231. (Parasite seed collected from Celtis, Ulmus, and Prosopis were grown on all three hosts. Growth was best on the host from which the seed was collected, suggesting some degree of host race development.)

Ehleringer, J R, Schulze, E D. 1985. Mineral concentration in an autoparasitic Phoradendron californicum growing on a parasitic P. californicum and its host Cercidium floridum. American Journal of Botany 72: 569-571. (Concentrations of a range of mineral elements were least in the host and highest in the hyperparasite.)

It is suggested this is due to higher transpiration rates in the parasite.)

Sahai, A, Shivanna, K R. 1985. Seed germination and seedling growth in Sopubia delphinifolia-a hemi-root parasite: germination requirements and requirements for seedling growth and the role of cotyledons. Annals of Botany 55: 775-783 and 785-791. (Light is shown to be essential for germination, which is also greatly increased chilling (4 C) for a few days or by ethylene. Continued growth in the absence of a host requires a carbohydrate source or high light intensity.)

Alosi, M C, Calvin, C L. 1985. The ultrastructure of dwarf mistletoe (Arceuthobium spp.) sinker cells in the region of the host secondary vasculature. Canadian Journal of Botany 63: 889-902. (Sinker cells are similar in three different species. Xylem is not continuous through the sinker cells but apoplastic continuity is provided by fused cellulosic cell walls.)

Musselman, L J. 1985. Bean stranglers! Explorer 27(3): 23-25. (A popular illustrated account of the genus Orobanch.)

Musselman, L J. 1985. Fertility and floral patterns in some species of Striga (Scrophulariaceae). National Geographic Society Research Reports 20: 487-491.

HAUSTORIUM is edited by L J Musselman, Department of Biological Sciences, Old Dominion University, Norfolk, Virginia 23508 USA and by C Parker, Tropical Weeds, Weed Research Organization, Begbroke Hill, Yarnton, Oxford OX5 1PF UK. Send material for publication in HAUSTORIUM as well as requests for copies to either editor.

This issue of HAUSTORIUM was reproduced and mailed to you by the Sorghum/Millet Collaborative Research Program, INTSORMIL. This collaborative research program is supported by a grant from the United States Agency for International Development, Contract Number DAN-1254-G-SS-5065-00.

